

Botrytis blight on Greenhouse Crops

Introduction

Botrytis blight is one of the most common diseases in the greenhouse. A wide range of plants including **greenhouse ornamentals, vegetables** and **herbs** are susceptible.

Favorable Conditions

Air currents and splashing water can easily disseminate the spores. In general, germination of spores and infection is dependent on a film of moisture for 8 to 12 hours, relative humidity of 93% or greater and temperatures between 55° and 65°F. Infection can occur through natural openings or wounds. After infection, colonization of plant tissues can occur at temperatures up to 70°F

Symptoms: *Botrytis* can cause leaf and flower spot and blights, stem cankers, damping off and cutting root rot. Plants may be attacked at any stage, but the new tender growth, and freshly injured tissues are most susceptible. *Botrytis* blight produces characteristic gray fuzzy appearing spores on the surface of infected tissues. However, spores only develop under humid conditions.

Figures 1 & 2: Botrytis sporulation and flowers with flecks of discoloration during humid conditions. Photos by L. Pundt



Leaf spots may develop a zonate pattern. Spots may develop where shedding flowers drop onto leaves below.

Figures 3 & 4: Leaf spots develop a zonate pattern. Photos by L. Pundt



Tan stem cankers can develop from blighted leaves on basil, fuchsia, poinsettia and other plants.

Figure 4 & 5: Tan stem cankers can develop. Photos by L. Pundt



Scouting

Concentrate scouting in areas where crop is closely spaced with poor air circulation, and on tender crops or more susceptible white flower crops. Look for dieback, tan stem cankers (especially near a wound), and gray fuzzy appearing spores during humid conditions.

Management

Management is challenging due to *Botrytis* abilities to survive as a saprophyte, rapidly invade host plants, and produce abundant spores that are easily moved within the greenhouse on air currents. *Botrytis* diseases can only be managed by a combination of methods including manipulation of environmental conditions (reducing humidity and duration of leaf wetness), sound cultural practices and proper use of fungicides. Fungicides alone cannot control *Botrytis* and this pathogen has a long history of fungicide resistance.

Follow Proper Sanitation Practices

- Control weeds and remove plant debris **before and during** production.
- Dispose of diseased plants and debris in a plastic trash bag. Keep the bag closed to help prevent spreading spores to uninfected plants as the bag is removed from the greenhouse.
- Cover trash cans to prevent the airborne spread of spores from diseased plant tissue.
- Avoid growing ornamental hanging baskets above especially susceptible crops or cultivars. The spent flowers dropping on plants below serve as an energy source for the fungus helping to encourage the development of *Botrytis* blight.

Follow Sound Cultural Practices

- Always water in the morning to reduce the length of time the leaves stay wet after irrigating to prevent foliar diseases. Rising temperatures during the day will evaporate water from the foliage, so the leaves stay dry.
- Avoid watering when the water will sit on leaf surfaces for long periods at the end of the day.
- Adequate spacing of plants and use of wire mesh benches help to improve air circulation.
- Proper plant nutrition and height management techniques help to prevent lush, overgrown plants thereby reducing humidity within the canopy.
- Avoid excess nitrogen fertility

- Calcium chloride has been shown to reduce susceptibility of some flowers.
- Follow proper planting and transplanting schedules to avoid holding plants any longer than necessary before their sale.

Specific Tips to Reduce Humidity levels in the Greenhouse

- Warm air holds more moisture than cool air. During warm days, the greenhouse air is more humid. As the air cools in the evening, the moisture-holding capacity drops until the dew point is reached. Water then begins to condense on surfaces.
- Reduce humidity by exhausting the moist air and replacing it with cooler outside air that is drier.

The method and time it takes to heat and vent depend upon the heating and ventilation system used in your greenhouse.

- In greenhouses with **vents**, turn the heat on and crack the vents open about one inch. The moist humid air escapes from the vents.
- In greenhouses with **fans**, activate the exhaust fans for a few minutes and then heat the greenhouse to raise the air temperature. Then, shut off the fans.
- A clock can be set to activate the fans. The cooler, outside air will lower humidity levels as it is warmed in the greenhouse.
- A relay may be needed to lock out the furnace or boiler until the fan shuts off so that flue gases are not drawn back into the greenhouse. (This will also help to prevent air pollution damage (ethylene or sulfur dioxide) to sensitive seedlings.)
- Heat and vent two or three times per hour in the evening after the sun goes down and early in the morning at sunrise.
- Heating and venting can be effective even if it is cool and raining outside.

Air movement, even in a closed greenhouse, helps reduce moisture on the plant surfaces and surrounding the plants. Using [horizontal airflow](#) (HAF) can also reduce condensation. HAF fans keep the air moving in the greenhouse, helping to minimize temperature differentials and cold spots where condensation occurs. Moving air is continually mixed. The mixed air along the surface does not cool below the dew point so does not condense on plant surfaces.

Selective Use of fungicides

Repeated applications of fungicides with the same mode of action (FRAC Group) can result in resistant populations of *Botrytis*. Widespread resistance to thiophanate-methyl (FRAC 1) has been reported from isolates collected in

1997 from Connecticut greenhouses. Isolates with resistance to iprodione (FRAC 2) were also collected but were not as common. Multiple resistance was also common. More recently (in 2013), resistance to fenhexamid (Decree) (FRAC 17) was reported from the Northeast.

Avoid making more than two consecutive applications with the same fungicide or products with the same mode of action. Product labels often have specific guidelines for resistance management.

The use of tank mixes of fungicide mixtures with different FRAC codes and rotations among different FRAC codes are critical for resistance management. [Biological fungicides](#) can also be part of a resistance management plan. However, they must be used **preventatively** for Botrytis management.

See the latest edition of the New England Greenhouse Floriculture Guide (A Management Guide for Insects, Diseases, Weeds and Growth Regulators) for the latest information on commercially available fungicides for Botrytis management on greenhouse ornamentals available from the [Northeast Greenhouse Conference & Expo](#).

At the end of the crop cycle, when crops are in flower, care must be exercised in selecting fungicides to avoid leaving unsightly residues or inadvertently damage the sensitive flowers. See [Tips on Preventing Botrytis during Cool, Cloudy Weather Periods](#) for more information.

For information on Botrytis Blight and Stem Canker on greenhouse tomatoes, see Battling Botrytis Blight and Stem Canker on Greenhouse Grown Tomatoes <http://www.e-gro.org/pdf/E408.pdf> and the latest edition of [the New England Vegetable Management Guide](#). <https://nevegetable.org/>

For more information on botrytis blight on herbs, see some selected fungicides labeled for use on herb bedding plants on the Greenhouse IPM website under herbs.

Botrytis blight is best management with proper environmental management combined with cultural controls. Use fungicides carefully as needed.

By Leanne Pundt, Extension Educator, UConn Extension, 2012, revised 2019

References

Chase, A.R., M.L. Daughtrey and R. A. Cloyd. 2018. Compendium of Bedding Plant Pests and Diseases. APS Press. 170 pp.

Daughtrey, M. L., R. L. Wick, and J. L. Peterson. 2000. Botrytis Blight of Flowering Potted Plants. Plant Health Progress. Reproduced from Compendium of Flowering Potted Plant Diseases. American Phytopathological Society. St. Paul, MN.

<http://www.plantmanagementnetwork.org/pub/php/management/botrytis/>

Hausbeck, M. 2017. B is for Botrytis blight on ornamentals. Michigan State Extension. https://www.canr.msu.edu/news/botrytis_blight_on_ornamentals

LaMondia, J. A. and S. M. Douglas. 1997. Sensitivity of *Botrytis cinerera* from Connecticut Greenhouses to Benzimidazole and Dicarboximide Fungicides. Plant Disease. 81:729-732.

Moorman, G. W. 2014. Botrytis or Gray Mold. Penn State Plant Disease Fact Sheet. <https://extension.psu.edu/botrytis-or-gray-mold>

Palmateer, A. 2017. Botrytis Blight on Greenhouse and Nursery Ornamentals. Grower Talks. <https://www.growertalks.com/Article/?articleid=23279>

Smith, T. S. and J.W. Bartok, Jr. 2004. Reduce humidity, Disease in Your Greenhouse. GMPro. November 2004, 52-58.

Trolinger, J. C. and D. L. Strider. 1985. *Botrytis* Diseases in Diseases of Floral Crops, Volume 1. D.L. Strider, Ed. Chapter 2.

Disclaimer for Fact Sheets:

The information in this document is for educational purposes only. The recommendations contained are based on the best available knowledge at the time of publication. Any reference to commercial products, trade or brand names is for information only, and no endorsement or approval is intended. UConn Extension does not guarantee or warrant the standard of any product referenced or imply approval of the product to the exclusion of others which also may be available. The University of Connecticut, UConn Extension, College of Agriculture, Health and Natural Resources is an equal opportunity program provider and employer.